

RINGS WHOSE TOTAL GRAPHS HAVE GENUS AT MOST ONE

HAMID REZA MAIMANI, CAMERON WICKHAM, AND SIAMAK YASSEMI

ABSTRACT. Let R be a commutative ring with $Z(R)$ its set of zero-divisors. In this paper, we study the total graph of R , denoted by $T(\Gamma(R))$. It is the (undirected) graph with all elements of R as vertices, and for distinct $x, y \in R$, the vertices x and y are adjacent if and only if $x + y \in Z(R)$. We investigate properties of the total graph of R and determine all isomorphism classes of finite commutative rings whose total graph has genus at most one (i.e., a planar or toroidal graph). In addition, it is shown that, given a positive integer g , there are only finitely many finite rings whose total graph has genus g .

INTRODUCTION

Let R be a commutative ring with non-zero unity. Let $Z(R)$ be the set of zero-divisors of R . The concept of the graph of the zero divisors of R was first introduced by Beck [6], where he was mainly interested in colorings. In his work all elements of the ring were vertices of the graph. This investigation of colorings of a commutative ring was then continued by D. D. Anderson and Naseer in [5]. In [4], D. F. Anderson and Livingston associate a graph, $\Gamma(R)$, to R with vertices $Z(R) \setminus \{0\}$, the set of non-zero zero-divisors of R , and for distinct $x, y \in Z(R) \setminus \{0\}$, the vertices x and y are adjacent if and only if $xy = 0$.

An interesting question was proposed by Anderson, Frazier, Lauve, and Livingston [3]: For which finite commutative rings R is $\Gamma(R)$ planar. A partial answer was given in [1], but the question remained open for local rings of order 32. In [12] and then independently in [7] and [13] it is shown that there is no ring of order 32 whose zero-divisor graph is planar.

The genus of a graph is the minimal integer n such that the graph can be drawn without crossing itself on a sphere with n handles (i.e. an oriented surface of genus n). Thus, a planar graph has genus zero, because it can be drawn on a sphere without self-crossing. In [13] and [16] the rings whose zero-divisor graph has genus one are studied. A genus one graph is called a toroidal graph. In other words, a graph G is toroidal if it can be embedded on the torus, that means, the graph's vertices can be placed on a torus such that no edges cross. Usually, it is assumed that G is also non-planar. In [17] it is shown that for a positive integer g , there are only finitely many finite rings whose zero-divisor graph has genus g .

In [2], D. F. Anderson and Badawi introduced the total graph of R , denoted by $T(\Gamma(R))$, as the graph with all elements of R as vertices, and for distinct $x, y \in R$, the vertices x and y are adjacent if and only if $x + y \in Z(R)$.

2000 *Mathematics Subject Classification.* 05C75, 13A15.

Key words and phrases. Total graph, genus, planar graph, toroidal graph.

The research of Hamid Maimani was in part supported by a grant from IPM (No. 88050214).

The research of Siamak Yassemi was in part supported by a grant from IPM (No. 88130213).

In this paper, we investigate properties of the total graph of R and determine all isomorphism classes of finite rings whose total graph has genus at most one (i.e., a planar or toroidal graph). In addition, we show that for a positive integer g , there are only finitely many finite rings whose total graph has genus g .

1. MAIN RESULT

A *complete graph* is a graph in which each pair of distinct vertices is joined by an edge. We denote the complete graph with n vertices by K_n . A *bipartite graph* is a graph such that its vertex set can be partitioned into two subsets V_1 and V_2 and each edge joins a vertex of V_1 to a vertex of V_2 . A *complete bipartite graph* is a bipartite graph such that each vertex in V_1 is joined by an edge to each vertex in V_2 , and is denoted by $K_{m,n}$ when $|V_1| = m$ and $|V_2| = n$. A *clique* of a graph is a maximal complete subgraph. For a graph G , the *degree* of a vertex v in G , denoted $\deg(v)$, is the number of edges of G incident with v . The number $\delta(G) = \min\{\deg(v) \mid v \text{ is a vertex of } G\}$ is the *minimum degree* of G . For a nonnegative integer k , a graph is called *k -regular* if every vertex has degree k . Recall that a graph is said to be *connected* if for each pair of distinct vertices v and w , there is a finite sequence of distinct vertices $v = v_1, \dots, v_n = w$ such that each pair $\{v_i, v_{i+1}\}$ is an edge. Such a sequence is said to be a *path* and the *distance* $d(v, w)$ between connected vertices v and w is the length of the shortest path connecting them. For any graph G , the disjoint union of k copies of G is denoted kG . Let S be a nonempty subset of vertex set of graph G . The *subgraph induced by S* is the subgraph with the vertex set S and with any edges whose endpoints are both in the S and is denoted by $\langle S \rangle$.

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two graphs with disjoint vertices set V_i and edges set E_i . The cartesian product of G_1 and G_2 is denoted by $G = G_1 \times G_2$ with vertices set $V_1 \times V_2$ and (x, y) is adjacent to (x', y') if $x = x'$ and y is adjacent to y' in G_2 or $y = y'$ and x is adjacent to x' in G_1 .

Lemma 1.1. *Let x be a vertex of $T(\Gamma(R))$. Then the degree of x is either $|Z(R)|$ or $|Z(R)| - 1$. In particular, if $2 \in Z(R)$, then $T(\Gamma(R))$ is a $(|Z(R)| - 1)$ -regular graph.*

Proof. If x is adjacent to y , then $x + y = a \in Z(R)$ and hence $y = a - x$ for some $a \in Z(R)$. We have two cases:

Case 1. Suppose $2x \in Z(R)$. Then x is adjacent to $a - x$ for any $a \in Z(R) \setminus \{2x\}$. Thus the degree of x is $|Z(R)| - 1$. In particular, if $2 \in Z(R)$, then $T(\Gamma(R))$ is a $(|Z(R)| - 1)$ -regular graph.

Case 2. Suppose $2x \notin Z(R)$. Then x is adjacent to $a - x$ for any $a \in Z(R)$. Thus the degree of x is $|Z(R)|$. $\square$

Let S_k denote the sphere with k handles, where k is a non-negative integer, that is, S_k is an oriented surface of genus k . The genus of a graph G , denoted by $\gamma(G)$, is the minimum integer n such that the graph can be embedded in S_n . A graph G is called a planar if $\gamma(G) = 0$, and toroidal if $\gamma(G) = 1$. We note here that if H is a subgraph of a graph G , then $\gamma(H) \leq \gamma(G)$.

In the following theorem we bring some well-known formulas, see, e.g., [14] and [15]:

Theorem 1.2. *The following statements hold:*

- (a) For $n \geq 3$ we have $\gamma(K_n) = \left\lceil \frac{(n-3)(n-4)}{12} \right\rceil$.
- (b) For $m, n \geq 2$ we have $\gamma(K_{m,n}) = \left\lceil \frac{(m-2)(n-2)}{4} \right\rceil$.
- (c) Let G_1 and G_2 be two graphs and for each i , p_i be the number of vertices of G_i . Then $\max\{p_1\gamma(G_2) + \gamma(G_1), p_2\gamma(G_1) + \gamma(G_2)\} \leq \gamma(G_1 \times G_2)$.

According to Theorem 1.2 we have $\gamma(K_n) = 0$ for $1 \leq n \leq 4$ and $\gamma(K_n) = 1$ for $5 \leq n \leq 7$ and for other value of n , $\gamma(K_n) \geq 2$.

Lemma 1.3. *Let $\mathbb{F}_q$ denote the field with q elements. Then the total graph of $\mathbb{F}_2 \times \mathbb{F}_q$ is isomorphic to $K_2 \times K_q$. Furthermore, for any positive integer m and $q > 2$,*

$$\gamma(T(\Gamma(\mathbb{F}_{2^m} \times \mathbb{F}_q))) \geq 2^m \left\lceil \frac{(q-3)(q-4)}{12} \right\rceil.$$

Proof. We induct on m . If $m = 1$, let $V_0 = \{(0, y) | y \in \mathbb{F}_q\}$ and let $V_1 = \{(1, y) | y \in \mathbb{F}_q\}$. Then the subgraphs G_i of the total graph of $\mathbb{F}_2 \times \mathbb{F}_q$ induced by each of the V_i is K_q . Now for each $y \in \mathbb{F}_q$, there is an edge between $(0, y) \in G_0$ and $(1, -y) \in G_1$. Furthermore, these are the only other edges in the total graph. Identifying $(1, -y)$ with $(1, y)$, we can replace G_1 with an isomorphic copy G'_1 ; under this isomorphism the edge between $(0, y) \in G_0$ and $(1, -y) \in G_1$ is the edge between $(0, y) \in G_0$ and $(1, y) \in G'_1$. Thus the total graph of $\mathbb{F}_2 \times \mathbb{F}_q$ has vertex set $\{(x, y) | x \in \mathbb{F}_2 \text{ and } y \in \mathbb{F}_q\}$, with an edge between (x, y) and (x', y') if $x = x'$ and $y \neq y'$, or $y = y'$ and $x \neq x'$. That is, it is the graph $K_2 \times K_q$. Parts (a) and (c) of Theorem 1.2 now yield

$$\gamma(T(\Gamma(\mathbb{F}_2 \times \mathbb{F}_q))) = \gamma(K_2 \times K_q) \geq 2 \left\lceil \frac{(q-3)(q-4)}{12} \right\rceil.$$

If $m > 1$, we can partition $\mathbb{F}_{2^m}$ into two sets, S_1 and S_2 , each of cardinality 2^{m-1} ; let f be a bijection from S_1 to S_2 . Since each element of a field of characteristic 2 is its own inverse, then the subgraph of $T(\Gamma(\mathbb{F}_{2^m} \times \mathbb{F}_q))$ induced by $S_i \times \mathbb{F}_q$ is isomorphic to $T(\Gamma(\mathbb{F}_{2^{m-1}} \times \mathbb{F}_q))$. For any $y \in \mathbb{F}_q$ and $s \in S_1$, the element (s, y) is adjacent to $(f(s), -y)$. We thus have a copy of $K_2 \times T(\Gamma(\mathbb{F}_{2^{m-1}} \times \mathbb{F}_q))$ as a subgraph of $T(\Gamma(\mathbb{F}_{2^m} \times \mathbb{F}_q))$. Part (c) of Theorem 1.2 and the induction hypothesis now yield

$$\begin{aligned} \gamma(T(\Gamma(\mathbb{F}_{2^m} \times \mathbb{F}_q))) &\geq \gamma(K_2 \times T(\Gamma(\mathbb{F}_{2^{m-1}} \times \mathbb{F}_q))) \\ &\geq 2\gamma(T(\Gamma(\mathbb{F}_{2^{m-1}} \times \mathbb{F}_q))) \\ &\geq 2^m \left\lceil \frac{(q-3)(q-4)}{12} \right\rceil. \end{aligned}$$

□

A *subdivision* of a graph is a graph obtained from it by replacing edges with pairwise internally disjoint paths. A remarkably simple characterization of planar graphs was given by Kuratowski in 1930. Kuratowski Theorem says that a graph is planar if and only if it contains no subdivision of K_5 or $K_{3,3}$ (see [2, p. 153]). In addition, every planar graph has a vertex v such that $\deg(v) \leq 5$.

Theorem 1.4. *For any positive integer g , There are finitely many finite rings R whose total graph has genus g .*

Proof. Let R be a finite ring. If R is local, then $Z(R)$ is the maximal ideal of R and $|R| \leq |Z(R)|^2$. If R is not local, then $R = R_1 \times R_2 \times \cdots \times R_n$ where each of the R_i 's is a local ring and $n \geq 2$ [10]. Suppose that $|R_1| \leq |R_2| \leq \cdots \leq |R_n|$ and set $R_1^* = 0 \times R_2 \times \cdots \times R_n$. Since $|R| = |R_1||R_1^*|$, we conclude that $|R| \leq |R_1^*|^2$. Let S denote either $Z(R)$ if R is local or R_1^* if R is not local. Then every pair of elements of S are adjacent in $T(\Gamma(R))$ and hence we have a complete graph $K_{|S|}$ in the structure of $T(\Gamma(R))$. This implies that $\gamma(K_{|S|}) \leq g$. Therefore $\lceil ((|S| - 3)(|S| - 4))/12 \rceil \leq g$ and so $|S| \leq \left(7 + \sqrt{49 + 48(g - 1)}\right)/2$, and hence $|R| \leq \left(\left(7 + \sqrt{49 + 48(g - 1)}\right)/2\right)^2$. $\square$

Theorem 1.5. *Let R be a finite ring such that $T(\Gamma(R))$ is planar. Then the following hold:*

- (a) *If R is local ring, then R is a field or R is isomorphic to the one of the 9 following rings:*

$$\mathbb{Z}_4, \mathbb{Z}_2[X]/(X^2), \mathbb{Z}_2[X]/(X^3), \mathbb{Z}_2[X, Y]/(X, Y)^2, \mathbb{Z}_4[X]/(2X, X^2), \\ \mathbb{Z}_4[X]/(2X, X^2 - 2), \mathbb{Z}_8, \mathbb{F}_4[X]/(X^2), \mathbb{Z}_4[X]/(X^2 + X + 1).$$

- (b) *If R is not local ring, then R is an infinite integral domain or R is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$ or $\mathbb{Z}_6$.*

Proof. Any planar graph has a vertex v with $\deg(v) \leq 5$. So if the total graph of R is planar, then $\delta(T(\Gamma(R))) \leq 5$. By Lemma 1.1, $\delta(T(\Gamma(R))) = |Z(R)|$ or $|Z(R)| - 1$, and hence $|Z(R)| \leq 6$.

(a) Assume that R is a local ring and let $n = |Z(R)|$ and $m = |R/Z(R)|$. If $2 \in Z(R)$ then $T(\Gamma(R)) \cong mK_n$ ([2, Theorems 2.1 and 2.2]). Hence $|Z(R)| \leq 4$. Also, $|R| = 2^k$ since $2 \in Z(R)$. So $|R| = 16, 8, 4$, or 2 . According to Corbas and Williams [8] there are two non-isomorphic rings of order 16 with maximal ideals of order 4, namely $\mathbb{F}_4[x]/(x^2)$ and $\mathbb{Z}_4[x]/(x^2 + x + 1)$ (see also Redmond [11]), so for these rings we have $T(\Gamma(R)) \cong 4K_4$. Since K_4 is planar we conclude that the total graphs of these rings are planar. In [8] it is also shown that there are 5 local rings of order 8 (except F_8). In all of these rings we have $|Z(R)| = 4$ and hence $T(\Gamma(R)) \cong 2K_4$. Also, there are two non-isomorphic local rings of order 4; these are $\mathbb{Z}_4$ and $\mathbb{Z}_2[X]/(X^2)$. For both we have $T(\Gamma(R)) \cong 2K_2$ and thus they are planar. Note that if $|Z(R)| = 1$, then R is a field and hence the total graph is planar. If $2 \notin Z(R)$, then $T(\Gamma(R)) \cong K_n \cup \left(\frac{m-1}{2}\right) K_{n,n}$ ([2, Theorem 2.2]). This implies $n \leq 2$, and thus R either has order 4 or is a field.

(b) Suppose that R is not local ring. Since R is finite, then there are finite local rings R_i such that $R = R_1 \times \cdots \times R_t$ where $t \geq 2$. Since $|Z(R)| \leq 6$ then we have the following candidates:

$$\mathbb{Z}_2 \times \mathbb{Z}_2, \mathbb{Z}_6, \mathbb{Z}_2 \times \mathbb{F}_4, \mathbb{Z}_2 \times \mathbb{Z}_4, \mathbb{Z}_2 \times \mathbb{Z}_2[X]/(X^2), \\ \mathbb{Z}_2 \times \mathbb{Z}_5, \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_3 \times \mathbb{F}_4.$$

The total graph of $\mathbb{Z}_2 \times \mathbb{Z}_2$ is isomorphic to the cycle C_4 , and this graph is planar. By Lemma 1.3, the total graph of $\mathbb{Z}_6 \cong \mathbb{Z}_2 \times \mathbb{Z}_3$ is isomorphic to $K_2 \times K_3$, which is also planar.

Let R be a ring with $|R| = n$. The subgraph of the total graph of $\mathbb{Z}_2 \times R$ induced by the set $\{0\} \times R$ is a copy of K_n . The edge $(1, 0) - (0, 0)$, together with the paths $(1, 0) - (1, -r) - (0, r)$ for each $r \in R$ yield a subdivision of K_{n+1} in the total

graph of $\mathbb{Z}_2 \times R$. Thus the total graphs of $\mathbb{Z}_2 \times \mathbb{F}_4$, $\mathbb{Z}_2 \times \mathbb{Z}_4$, $\mathbb{Z}_2 \times \mathbb{Z}_2[X]/(X^2)$, and $\mathbb{Z}_2 \times \mathbb{Z}_5$ are not planar. Also, the total graph of $\mathbb{Z}_3 \times R$ contains a subgraph which is isomorphic to $K_{3,n}$ (consider the induced subgraph $\langle S \rangle$ where $S = \{(1, r) \mid r \in R\} \cup \{(2, r) \mid r \in R\}$). Thus the total graphs of $\mathbb{Z}_3 \times \mathbb{Z}_3$ and $\mathbb{Z}_3 \times \mathbb{F}_4$ are not planar. $\square$

Theorem 1.6. *Let R be a finite ring such that $T(\Gamma(R))$ is toroidal. Then the following statements hold:*

- (a) *If R is local ring, then R is isomorphic to $\mathbb{Z}_9$, or $\mathbb{Z}_3[X]/(X^2)$.*
- (b) *If R is not local ring, then R is isomorphic to one of the following rings:*

$$\mathbb{Z}_2 \times \mathbb{F}_4, \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_2 \times \mathbb{Z}_4, \mathbb{Z}_2 \times \mathbb{Z}_2[x]/(x^2), \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$$

Proof. For any graph G with ν vertices and genus g we have $\delta(G) \leq 6 + (12g - 12)/\nu$. If $\gamma(G) = 1$, then $\delta(G) \leq 6$ and equality holds if and only if G is a triangulation of the torus and 6-regular (see [16, Proposition 2.1]). If R is a finite ring with toroidal total graph, then $\delta(T(\Gamma(R))) \leq 6$, and by Lemma 1.1 $\delta(T(\Gamma(R))) = |Z(R)|$ or $|Z(R)| - 1$. Thus we conclude that $|Z(R)| \leq 7$.

(a) Let R be a local ring. If $2 \in Z(R)$, then $T(\Gamma(R))$ is a disjoint union of copies of the complete graph K_n , where $|Z(R)| = n$. Hence $5 \leq n \leq 7$. But in this case $|Z(R)|$ is a power of 2 and thus there are no such local rings. Now suppose that $2 \notin Z(R)$. Then $T(\Gamma(R)) \cong K_n \cup (\frac{m-1}{2}) K_{n,n}$, where $n = |Z(R)|$ and $m = |R/Z(R)|$. Thus $3 \leq n \leq 4$ and since $2 \notin Z(R)$ we must have $n = 3$. There are two local rings, $\mathbb{Z}_9$ and $\mathbb{Z}_3[X]/(X^2)$, such that the cardinality of the set of the zero-divisors is 3; both of these rings have total graph $K_3 \cup K_{3,3}$ which is toroidal.

(b) Assume that R is not a local ring. Since $|Z(R)| \leq 7$, we have the following candidates for R by Theorem 1.5(b):

$$\mathbb{Z}_2 \times \mathbb{F}_4, \mathbb{Z}_2 \times \mathbb{Z}_4, \mathbb{Z}_2 \times \mathbb{Z}_2[x]/(x^2), \mathbb{Z}_2 \times \mathbb{Z}_5, \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_3 \times \mathbb{F}_4, \mathbb{F}_4 \times \mathbb{F}_4, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2.$$





By Theorem 1.5, $\gamma(T(\Gamma(\mathbb{Z}_2 \times \mathbb{F}_4)))$, $\gamma(T(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_4)))$, $\gamma(T(\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_3)))$ are all at least 1. The embeddings in Figure 1 parts (a), (b), and (c) show explicitly that $\gamma(T(\Gamma(\mathbb{Z}_2 \times \mathbb{F}_4))) = \gamma(T(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_4))) = \gamma(T(\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_3))) = 1$. Since $T(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2[x]/(x^2))) \cong T(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_4))$, then $\gamma(T(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2[x]/(x^2)))) = 1$.

If we partition the elements of $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ by the four sets $V_1 = \{(0, 0, 0), (1, 1, 1)\}$, $V_2 = \{(1, 0, 0), (0, 1, 1)\}$, $V_3 = \{(0, 1, 0), (1, 0, 1)\}$, and $V_4 = \{(0, 0, 1), (1, 1, 0)\}$, it is clear that $T(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2)) = K_{2,2,2,2}$. Hence $\gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2) = 1$ by [9, Corollary 4]

By Lemma 1.3, $\gamma(T(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_5))) \geq 2$ and thus is not toroidal. By (the proof of) Lemma 1.3, $\gamma(T(\Gamma(\mathbb{F}_4 \times \mathbb{F}_4))) \geq 2\gamma(T(\Gamma(\mathbb{Z}_2 \times \mathbb{F}_4)))$ and so by Theorem 1.5 $T(\Gamma(\mathbb{F}_4 \times \mathbb{F}_4))$ is not toroidal.

Finally, Figure 1 part (d) shows a subgraph of $T(\Gamma(\mathbb{Z}_3 \times \mathbb{F}_4))$ that is a subdivision of $K_{5,4}$, and thus by Theorem 1.2 $\gamma(T(\Gamma(\mathbb{Z}_3 \times \mathbb{F}_4))) \geq 2$. $\square$

The authors wish to thank the referee for the detailed and useful comments.

FIGURE 1. Embeddings in the torus and a subgraph of $T(\Gamma(\mathbb{Z}_3 \times \mathbb{F}_4))$ (a) Embedding of $\mathbb{Z}_2 \times \mathbb{Z}_4$ (b) Embedding of $\mathbb{Z}_2 \times \mathbb{F}_4$ (c) Embedding of $\mathbb{Z}_3 \times \mathbb{Z}_3$ (d) A subgraph of $T(\Gamma(\mathbb{Z}_3 \times \mathbb{F}_4))$
that is a subdivision of $K_{5,4}$

REFERENCES

1. S. Akbari, H. R. Maimani and S. Yassemi, *When a zero-divisor graph is planar or a complete r -partite graph*, J. Algebra **270** (2003), 169–180.
2. D. F. Anderson and A. Badawi, *The total graph of a commutative ring*, J. Algebra **320** (2008), 2706–2719.
3. D. F. Anderson, A. Frazier, A. Lauve, and P. S. Livingston, *The zero-divisor graph of a commutative ring, II*, In *Ideal theoretic methods in commutative algebra (Columbia, MO, 1999)*, 61–72, Dekker, New York, 2001.
4. D. F. Anderson and P. S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999), 434–447.
5. D. D. Anderson and M. Naseer, *Beck's coloring of a commutative ring*, J. Algebra **159** (1993), 500–514.
6. I. Beck, *Coloring of commutative rings*, J. Algebra **116** (1988), no. 1, 208–226.
7. R. Belshoff, J. Chapman, *Planar zero-divisor graphs*, J. Algebra **316** (2007), 471–480.
8. B. Corbas, G.D. Williams, *Ring of order p^5 . II. Local rings*, J. Algebra **231**(2) (2000) 691–704.
9. D. L. Craft, *On the genus of joins and composition of graphs*, Discrete Math. **178** (1998) 25–50.
10. B. R. McDonald, *Finite Rings with Identity*, Pure and Applied Mathematics, vol. 28, Marcel Dekker, Inc., New York, 1974.
11. S.P. Redmond, *On zero-divisor graphs of small finite commutative rings*, Discrete Math. **307** (2007), no. 9–10, 1155–1166.
12. Neal O. Smith, *Planar zero-divisor graphs*, Internat. J. Commutative Rings **2** (2002), 177–188.
13. H. Wang, *Zero-divisor graphs of genus one*, J. Algebra **304** (2006), 666–678.
14. A. T. White, *Graphs, Groups and Surfaces*, North-Holland, Amsterdam, 1973.
15. A. T. White, *The genus of the Cartesian product of two graphs*, J. Combin. Theory Ser. B **11** (1971), 89–94.
16. C. Wickham, *Classification of rings with genus one zero-divisor graphs*, Comm. Algebra **36** (2008), 325–345.
17. C. Wickham, *Rings whose zero-divisor graphs have positive genus*, J. Algebra **321** 2009, 377–383.

HAMID REZA MAIMANI, DEPARTMENT OF MATHEMATICS, SHAHID RAJAEI UNIVERSITY, TEHRAN, IRAN, AND INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), TEHRAN IRAN.

CAMERON WICKHAM, MATHEMATICS DEPARTMENT, MISSOURI STATE UNIVERSITY, SPRINGFIELD, MO 65897, USA.

SIAMAK YASSEMI, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEHRAN, TEHRAN, IRAN, AND INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), TEHRAN IRAN.